


AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1 (currently amended). A compound of formula I,



wherein

R¹ represents C₁₋₁₂ alkyl, C₃₋₁₂ cycloalkyl, -(CH₂)_a-aryl, or (CH₂)_aHet¹ (all of which are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C₁₋₄ alkyl, C₃₋₄ cycloalkyl and/or C₁₋₄ alkoxy or C₃₋₄ cycloalkoxy);

a represents 0, 1, 2, 3, or 4;

Het¹ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

X represents O or S;

R^{5a} and R^{5b} independently represent H, C₁₋₃ alkyl or C₃ cycloalkoxy;

R² and R³ independently represent H, C₁₋₄ alkyl (optionally substituted with one or more nitro or cyano groups), C₃₋₄ cycloalkyl, OR⁷, N(R^{7a})R^{7b}, OC(O)R⁸ or together form -O-(CH₂)₂-O-, -(CH₂)₃-, -(CH₂)₄- or -(CH₂)₅-;

R⁷ and R⁸ independently represent H, C₁₋₆ alkyl, or -(CH₂)_b-aryl (which latter two groups are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C₁₋₄ alkyl, C₁₋₄ alkoxy, and/or C₃₋₄ cycloalkyl);

R^{7a} and R^{7b} independently represent H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

b represents 0, 1, 2, 3 or 4;

R⁴ represents H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

D represents H, -OH, or -(CH₂)_cN(R¹⁰)(R¹¹);

c represents 0, 1, 2, 3 or 4;

R¹⁰ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, -(CH₂)_d-aryl, -C(NH)NH₂, -S(O)₂R¹³, -[C(O)]_eN(R¹⁴)(R¹⁵), -C(O)R¹⁶ or -C(O)OR¹⁷;

e represents 1 or 2;

R¹¹ represents H, C₁₋₆ alkyl, -C(O)R¹⁸ or -(CH₂)_f-aryl (which latter group is optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkyl and/or C₃₋₆ cycloalkoxy);

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} independently represent H, C_{1-6} alkyl, C_{3-6} cycloalkyl, Het^2 or $-(CH_2)_g$ -aryl (which latter three groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{13} represents C_{1-6} alkyl, C_{3-6} cycloalkyl, aryl or $-(CH_2)_h$ -aryl (all of which are all optionally substituted by one or more substituents chosen from halo, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

d, f, g and h independently represent 0, 1, 2, 3 or 4;

Het^2 represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

R^6 represents one or more optional substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{18a}$), C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{3-6} cycloalkoxy, $-C(O)N(H)R^{19}$, $-NHC(O)N(H)R^{20}$, $-N(H)S(O)_2R^{21}$ and/or $-OS(O)_2R^{22}$;

R^{19} and R^{20} independently represent H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

R^{18a} , R^{21} and R^{22} independently represent C_{1-6} alkyl or C_{3-6} cycloalkyl;

A represents a single bond, C_{1-6} alkylene, $-N(R^{23})(CH_2)_j$ -, $-O(CH_2)_j$ - or $-(CH_2)_jC(H)(OR^{23})(CH_2)_k$ - $-(CH_2)_iC(H)(OR^{23})(CH_2)_k$ - (in which latter three groups, the $-(CH_2)_j$ - group is attached to the bispidine nitrogen atom, and which latter four groups are all optionally substituted by one or more OH groups);

B represents a single bond, C_{1-4} alkylene, $-(CH_2)_mN(R^{24})$ -, $(CH_2)_mS(O)_n$ -, $-(CH_2)_mO$ - (in which three latter groups, the $-(CH_2)_m$ - group is attached to the carbon

atom bearing D and R⁴), -C(O)N(R²⁴)- (in which latter group, the -C(O)- group is attached to the carbon atom bearing D and R⁴), N(R²⁴)C(O)O(CH₂)_m- or -N(R²⁴)(CH₂)_m- (in which latter two groups, the N(R²⁴) group is attached to the carbon atom bearing D and R⁴);

j, k and m independently represent 0, 1, 2, 3 or 4;

n represents 0, 1 or 2;

R²³ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or C(O)R²⁵

R²⁴ represents H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

R²⁵ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Het³ or -(CH₂)_p-aryl (which latter two groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkyl and/or C₃₋₆ cycloalkoxy);

Het³ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

p represents 0, 1, 2, 3 or 4;

or a pharmaceutically acceptable salt, N-oxide or C₁₋₄ alkyl quaternary ammonium derivative thereof;

wherein alkyl groups that R¹, R², R³, R⁴, R^{5a}, R^{5b}, R⁶, R⁷, R^{7a}, R^{7b}, R⁸, R¹⁰, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R^{18a}, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and D may represent, and with which R¹, R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted; and alkoxy groups and that R⁶ may represent, and with which R¹, R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted, may be linear or, when there is a sufficient number (i.e. three) of carbon atoms, be branched and/or cyclic cycloalkyl or

cycloalkyl with carbon ranges as defined above, and wherein, when there is a sufficient number (i.e. four) of carbon atoms, such alkyl and alkoxy groups may also be part ~~cyclic/acyclic~~ cycloalkyl/acyclic or cycloalkoxy/acyclic, with carbon ranges as defined above, and wherein such alkyl and alkoxy groups may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen and/or substituted by one or more fluoro groups; and

wherein alkylene groups that A and B may represent, and $-(CH_2)-$ containing groups that R^1 , R^2 and R^3 (together), R^7 , R^8 , R^{10} , R^{11} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{25} , A, B and D may include, may be linear or, when there is a sufficient number (i.e. two) of carbon atoms, be branched, and wherein such alkylene groups and $-(CH_2)-$ containing chains may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen;

provided that:

(a) when D represents either H or -OH, and R^{5a} and R^{5b} both represent H, then at least one of R^2 and R^3 represents OR^7 , $OC(O)R^8$ or C_{1-4} alkyl, which alkyl group is substituted with one or more nitro or cyano groups; and

(b) when D represents -OH or $-(CH_2)_cN(R^{10})R^{11}$ in which c represents 0, then:-

(i) A does not represent $-N(R^{23})(CH_2)_j-$, $-O(CH_2)_j-$ or $-CH_2)_jC(H)(OR^{23})(CH_2)_k-$ (in which k is 0); and/or

(ii) m does not represent 0 when B represents $-(CH_2)_mN(R^{24})-$, $-(CH_2)_mS(O)_n-$ or $-(CH_2)_mO-$.

2 (currently amended). A compound as claimed in Claim 1, wherein R¹ represents optionally substituted -(CH₂)_a-phenyl, in which a is 0, 1, 2 or 3, or optionally substituted, optionally unsaturated, linear, branched ~~or cyclic~~, C₁₋₁₈ alkyl or C₃₋₁₈ cycloalkyl (which latter C₁₋₁₈ alkyl or C₃₋₁₈ cycloalkyl group may also be interrupted by an oxygen atom).

3 (previously presented). A compound as claimed in Claim 1, wherein R² represents H, OR⁷, -CH₂NO₂ or -OC(O)R⁸ or together with R³ -O-(CH₂)₂-O-.

4 (previously presented). A compound as claimed in Claim 1, wherein R³ represents H, OR⁷, C₁₋₄ alkyl or together with R² represents -O-(CH₂)₂-O-.

5 (previously presented). A compound as claimed in Claim 1, wherein R⁴ represents H or C₁₋₂ alkyl.

6 (previously presented). A compound as claimed in Claim 1, wherein R^{5a} and R^{5b} either both represent H or both represent methyl.

7 (previously presented). A compound as claimed in Claim 1, wherein R⁶ represents one or more substituents selected from C₁₋₆ alkyl, cyano, nitro, amino or C(O)N(H)R¹⁹ or N(H)S(O)₂R²¹.

8 (previously presented). A compound as claimed in Claim 1, wherein X represents O.

9 (previously presented). A compound as claimed in Claim 1, wherein A represents a single bond or linear, or branched, C₁₋₄ alkylene (which group is also optionally interrupted by O).

10 (previously presented). A compound as claimed in Claim 1, wherein B represents a single bond, C₁₋₄ alkylene, -(CH₂)_mO- or -(CH₂)_mN(R²⁴)- (in which latter two cases m is 1, 2 or 3).

11 (previously presented). A compound as claimed in Claim 1, wherein when D represents -(CH₂)_cN(R¹⁰)(R¹¹), c represents 0, 1 or 2.

12 (previously presented). A compound as claimed in Claim 1, wherein when D represents -(CH₂)_cN(R¹⁰)(R¹¹), R¹⁰ represents H, C₁₋₄ alkyl, -C(O)R¹⁶ (in which R¹⁶ is H, C₁₋₃ alkyl or Het²), -C(O)OR¹⁷ (in which R¹⁷ is C₁₋₅ alkyl, phenyl or C₁₋₃ alkylphenyl), -C(NH)NH₂ or [C(O)]₆N(H)R₁₅ (in which R₁₅ is H or C₁₋₃ alkyl).

13 (previously presented). A compound as claimed in Claim 1, wherein when D represents -(CH)_cN(R¹⁰)(R¹¹), R¹¹ represents H.


14 (previously presented). A pharmaceutical formulation including a compound as defined in Claim 1 in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

15-19 (cancelled).

20 (previously presented). A method of prophylaxis or treatment of an arrhythmia which method comprises administration of a therapeutically effective amount of a compound as defined in Claim 1 to a person in need thereof.

21 (previously presented) A process for the preparation of a compound of formula I as defined in Claim 1 which comprises:

(a) reaction of a compound of formula II,




II

wherein R^2 , R^3 , R^4 , R^{5a} , R^{5b} , R^6 , A, B and D are as defined in Claim 1 with a compound of formula III,




wherein L^1 represents a leaving group and R^1 and X are as defined in Claim 1;

(b) for compounds of formula I in which A represents CH_2 and D represents $-OH$ or $N(R^{10})H$, reaction of a compound of formula IV,




wherein R^1 , R^2 , R^3 , R^{5a} , R^{5b} and X are as defined in Claim 1, with a compound of formula V,



wherein Y represents O or $N(R^{10})$ and R^4 , R^6 , R^{10} and B are as defined in Claim


1;

(c) reaction of a compound of formula IV, as defined above, with a compound of formula VI,




wherein L² represents a leaving group and R⁴, R⁶, A, B and D are as defined in Claim 1;

(d) for compounds of formula I in which D represents H or OH and R⁴ represents H, reduction of a compound of formula VII,



wherein R^1 , R^2 , R^3 , R^{5a} , R^{5b} , R^6 , A, B and X are as defined in Claim 1;

(e) for compounds of formula I in which one of R^2 and R^3 represents H or OH and the other represents H, reduction of a corresponding compound of formula VIII,



wherein R^1 , R^4 , R^{5a} , R^{5b} , R^6 , A, B, D and X are as defined in Claim 1;

(f) for compounds of formula I in which R^2 and/or R^3 represents $OC(O)R^8$ and R^8


is as defined in Claim 1, coupling of a corresponding compound of formula I in which R^2 and/or R^3 (as appropriate) represents OH and a compound of formula VIIIA,



VIIIA

wherein R^8 is as defined in Claim 1;

(g) for compounds of formula I in which D represents $-(CH_2)_cNH_2$, reduction of a corresponding compound of formula IX,



wherein c , R^1 , R^2 , R^3 , R^4 , R^{5a} , R^{5b} , R^6 , A, B and X are as defined in Claim 1;

(h) for compounds of formula I in which D represents $-N(R^{11})C(O)NH(R^{15})$, in which R^{11} and R^{15} are as defined in Claim 1 except that R^{11} does not represent $C(O)R^{18}$, reaction of a corresponding compound of formula I in which D represents $-N(R^{11})H$, in which R^{11} is as defined in Claim 1 except that it does not represent $C(O)R^{18}$ in which R^{18} is as defined in Claim 1, with a compound of formula X,



wherein R^{15} is as defined in Claim 1;

(i) for compounds of formula I in which D represents $-\text{N}(\text{H})[\text{C}(\text{O})]_2\text{NH}_2$,

reaction of a corresponding compound of formula I in which D represents

$-\text{NH}_2$ with oxalic acid diamide;

(j) for compounds of formula I in which D represents $-\text{N}(\text{R}^{11})\text{C}(\text{O})\text{R}^{16}$, in

which R^{11} and R^{16} are as defined in Claim 1 except that R^{11} does not represent $\text{C}(\text{O})\text{R}^{18}$,

reaction of a corresponding compound of formula I in which D represents $-\text{N}(\text{R}^{11})\text{H}$, in

which R^{11} is as defined in Claim 1 except that it does not represent $\text{C}(\text{O})\text{R}^{18}$ in which

R^{18} is as defined in Claim 1, with a compound of formula XI,



wherein R_x represents a suitable leaving group and R^{16} is as defined in Claim 1;

(k) for compounds of formula I in which D represents $-\text{N}(\text{H})\text{R}^{10}$ and R^{10} is as

defined in Claim 1 except that it does not represent H or $-\text{C}(\text{NH})\text{NH}_2$, reaction of a

corresponding compound of formula I wherein D represents $-\text{NH}_2$ with a compound of formula XIA,



wherein R^{10a} represents R^{10} as defined in Claim 1 except that it does not represent H or $-\text{C}(\text{NH})\text{NH}_2$ and L^1 is as defined above;

(l) for compounds of formula I which are bispidine-nitrogen N-oxide derivatives, oxidation of the corresponding bispidine nitrogen of a corresponding compound of formula I;

(m) for compounds of formula I which are C_{1-4} alkyl quaternary ammonium salt


derivatives, in which the alkyl group is attached to a bispidine nitrogen, reaction, at the bispidine nitrogen, of a corresponding compound of formula I with a compound of formula XII,



XII


wherein R^a represents C_{1-4} alkyl and Hal represents Cl, Br or I;

(n) for compounds of formula I in which D and R^4 both represent H, A represents C_{1-6} alkylene, B represents $N(R^{24})(CH_2)_m$ and m and R^{24} are as defined in Claim 1, reaction of a compound of formula XIII,



XIII

wherein A^a represents C_{1-6} alkylene and R^1 , R^2 , R^3 , R^{5a} , R^{5b} , R^{24} and X are as defined in Claim 1 with a compound of formula XIV,



XIV


wherein R^6 , m are as defined in Claim 1 and Hal is as defined above;

(o) reaction of a compound of formula II, as defined above, with a compound of formula XV,



wherein R^1 and X are as defined in Claim 1, in the presence of 1,1'-carbonyldiimidazole;

(p) for compounds of formula I in which one of R^2 and R^3 represents $—NH_2$ and the other represents H, reduction of a compound of formula XVA,




wherein R^1 , R^4 , R^{5a} , R^{5b} , R^6 , A, B, D and X are as defined in Claim 1; or

(q) for compounds of formula I in which one or both of R^2 and R^3 represent $-N(R^{7a})R^{7b}$ in which one or both of R^{7a} and R^{7b} represents C_{1-6} alkyl, alkylation of a corresponding compound of formula I in which R^2 and/or R^3 represent $-N(R^{7a})R^{7b}$ (as appropriate) in which R^{7a} and/or R^{7b} (as appropriate) represent H, using a compound of formula XXIB,



wherein R^{7c} represents C_{1-6} alkyl and L^1 is as defined above.

22 (currently amended). A compound of formula II



wherein R^{5a} and R^{5b} independently represent H, C_{1-3} alkyl or C_3 cycloalkoxy;

R^2 and R^3 independently represent H, C_{1-4} alkyl (optionally substituted with one or more nitro or cyano groups), C_{3-4} cycloalkyl, OR^7 , $N(R^{7a})R^{7b}$, $OC(O)R^8$ or together form -
 $O-(CH_2)_2-O-$, $-(CH_2)_3-$, $-(CH_2)_4-$ or $-(CH_2)_5-$;

R^7 and R^8 independently represent H, C_{1-6} alkyl, or $-(CH_2)_b$ -aryl (which latter two groups are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, and/or C_{3-4} cycloalkyl);

R^{7a} and R^{7b} independently represent H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

b represents 0, 1, 2, 3 or 4;

R^4 represents H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

D represents H, -OH, or $-(CH_2)_cN(R^{10})(R^{11})$;

c represents 0, 1, 2, 3 or 4;

R^{10} represents H, C_{1-6} alkyl, C_{3-6} cycloalkyl, $-(CH_2)_d$ -aryl, $-C(NH)NH_2$, $-S(O)_2R^{13}$, $-[C(O)]_eN(R^{14})(R^{15})$, $-C(O)R^{16}$ or $-C(O)OR^{17}$;

e represents 1 or 2;

R^{11} represents H, C_{1-6} alkyl, $-C(O)R^{18}$ or $-(CH_2)_f$ -aryl (which latter group is optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} independently represent H, C_{1-6} alkyl, C_{3-6} cycloalkyl, Het² or $-(CH_2)_g$ -aryl (which latter three groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{13} represents C_{1-6} alkyl, C_{3-6} cycloalkyl, aryl or $-(CH_2)_h$ -aryl (all of which are optionally substituted by one or more substituents chosen from halo, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

d, f, g and h independently represent 0, 1, 2, 3 or 4;

Het² represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

R^6 represents one or more optional substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{18a}$), C_{1-6} alkoxy, C_{3-6}

cycloalkyl, C₃₋₆ cycloalkoxy, -C(O)N(H)R¹⁹, -NHC(O)N(H)R²⁰, -N(H)S(O)₂R²¹ and/or -OS(O)₂R²²;

R¹⁹ and R²⁰ independently represent H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

R^{18a}, R²¹ and R²² independently represent C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

A represents a single bond, C₁₋₆ alkylene, -N(R²³)(CH₂)_j-, -O(CH₂)_j- or

-(CH₂)_jC(H)(OR²³)(CH₂)_k- -(CH₂)_jC(H)(OR²³)(CH₂)_k- (in which latter three groups, the -
 (CH₂)_j- group is attached to the bispidine nitrogen atom, and which latter four groups are
 all optionally substituted by one or more OH groups);

B represents a single bond, C₁₋₄ alkylene, -(CH₂)_mN(R²⁴)-, (CH₂)_mS(O)_n-,
 -(CH₂)_mO- (in which three latter groups, the -(CH₂)_m- group is attached to the carbon
 atom bearing D and R⁴), -C(O)N(R²⁴)- (in which latter group, the -C(O)- group is
 attached to the carbon atom bearing D and R⁴), N(R²⁴)C(O)O(CH₂)_m- or -N(R²⁴)(CH₂)_m-
 (in which latter two groups, the N(R²⁴) group is attached to the carbon atom bearing D
 and R⁴);

j, k and m independently represent 0, 1, 2, 3 or 4;

n represents 0, 1 or 2;

R²³ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or C(O)R²⁵

R²⁴ represents H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

R²⁵ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Het³ or -(CH₂)_p-aryl (which latter two
 groups are optionally substituted by one or more substituents selected from -OH, cyano,
 halo, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkyl and/or C₃₋₆ cycloalkoxy);

Het³ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;


p represents 0, 1, 2, 3 or 4;

wherein alkyl groups that R², R³, R⁴, R^{5a}, R^{5b}, R⁶, R⁷, R^{7a}, R^{7b}, R⁸, R¹⁰, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R^{18a}, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and D may represent, and with which R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted; and alkoxy groups and that R⁶ may represent, and with which R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted, may be linear or, when there is a sufficient number (i.e. three) of carbon atoms, be branched and/or ~~cyclic~~ cycloalkyl or cycloalkoxy with carbon ranges as defined above, and wherein, when there is a sufficient number (i.e. four) of carbon atoms, such alkyl and alkoxy groups may also be part ~~cyclic/acyclic~~ cycloalkyl/acyclic or cycloalkoxy/acyclic with carbon ranges as defined above, and wherein such alkyl and alkoxy groups may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen and/or substituted by one or more fluoro groups; and

wherein alkylene groups that A and B may represent, and -(CH₂)- containing groups that R² and R³ (together), R⁷, R⁸, R¹⁰, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R²⁵, A, B and D may include, may be linear or, when there is a sufficient number (i.e. two) of carbon atoms, be branched, and wherein such alkylene groups and -(CH₂)- containing chains may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen,

provided that when R^{5a} and R^{5b} both represent H, then D does not represent H or OH.

23 (currently amended). A compound of formula IV



wherein R^1 represents C_{1-12} alkyl, C_{3-12} cycloalkyl, $-(CH_2)_a$ -aryl, or $(CH_2)_a$ Het¹ (all of which are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{3-4} cycloalkyl and/or C_{1-4} alkoxy or C_{3-4} cycloalkoxy);

a represents 0, 1, 2, 3, or 4;

Het¹ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

X represents O or S;

R^{5a} and R^{5b} independently represent H, C_{1-3} alkyl or C_3 cycloalkoxy;

R^2 and R^3 independently represent H, C_{1-4} alkyl (optionally substituted with one or more nitro or cyano groups), C_{3-4} cycloalkyl, OR^7 , $N(R^{7a})R^{7b}$, $OC(O)R^8$ or together form -
 $O-(CH_2)_2-O-$, $-(CH_2)_3-$, $-(CH_2)_4-$ or $-(CH_2)_5-$;

R^7 and R^8 independently represent H, C_{1-6} alkyl, or $-(CH_2)_b$ -aryl or (which latter two groups are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{1-4} alkoxy, and/or C_{3-4} cycloalkyl);


R^{7a} and R^{7b} independently represent H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

b represents 0, 1, 2, 3 or 4;

wherein alkyl groups that R^1 , R^2 , R^3 , R^{5a} , R^{5b} , R^7 , R^{7a} , R^{7b} and R^8 may represent, and with which R^1 , R^7 and R^8 may be substituted; and alkoxy groups and with which R^1 , R^7 and R^8 may be substituted, may be linear or, when there is a sufficient number (i.e. three) of carbon atoms, be branched and/or ~~cyclic~~ cycloalkyl or cycloalkoxy with carbon ranges as defined above, and wherein, when there is a sufficient number (i.e. four) of carbon atoms, such alkyl and alkoxy groups may also be part ~~cyclic/acyclic~~ cycloalkyl/acyclic or cycloalkoxy/acyclic with carbon ranges as defined above, and wherein such alkyl and alkoxy groups may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen and/or substituted by one or more fluoro groups;

provided that when R^{5a} and R^{5b} both represent H, then at least one of R^2 and R^3 represents OR^7 , $OC(O)R^8$ or C_{1-4} alkyl, which alkyl group is substituted with one or more nitro or cyano groups.

24 (currently amended). A compound of formula VIII



wherein R^1 represents C_{1-12} alkyl, C_{3-12} cycloalkyl, $-(CH_2)_a$ -aryl, or $(CH_2)_a$ Het¹ (all of which are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{3-4} cycloalkyl and/or C_{1-4} alkoxy or C_{3-4} cycloalkoxy);

a represents 0, 1, 2, 3, or 4;

Het¹ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

X represents O or S;

R^{5a} and R^{5b} independently represent H, C_{1-3} alkyl or C_3 cycloalkoxy;

R^4 represents H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

D represents H, -OH, or $-(CH_2)_cN(R^{10})(R^{11})$;

c represents 0, 1, 2, 3 or 4;

R^{10} represents H, C_{1-6} alkyl, C_{3-6} cycloalkyl, $-(CH_2)_d$ -aryl, $-C(NH)NH_2$, $-S(O)_2R^{13}$, $-[C(O)]_eN(R^{14})(R^{15})$, $-C(O)R^{16}$ or $-C(O)OR^{17}$;

e represents 1 or 2;

R^{11} represents H, C_{1-6} alkyl, $-C(O)R^{18}$ or $-(CH_2)_f$ -aryl (which latter group is optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} independently represent H, C_{1-6} alkyl, C_{3-6} cycloalkyl, Het² or $-(CH_2)_g$ -aryl (which latter three groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{13} represents C_{1-6} alkyl, C_{3-6} cycloalkyl, aryl or $-(CH_2)_h$ -aryl (all of which are all optionally substituted by one or more substituents chosen from halo, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

d, f, g and h independently represent 0, 1, 2, 3 or 4;

Het² represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

R^6 represents one or more optional substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{18a}$), C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{3-6} cycloalkoxy, $-C(O)N(H)R^{19}$, $-NHC(O)N(H)R^{20}$, $-N(H)S(O)_2R^{21}$ and/or $-OS(O)_2R^{22}$;

R^{19} and R^{20} independently represent H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

R^{18a} , R^{21} and R^{22} independently represent C_{1-6} alkyl or C_{3-6} cycloalkyl;

A represents a single bond, C₁₋₆ alkylene, -N(R²³)(CH₂)_j-, -O(CH₂)_j- or -(CH₂)_jC(H)(OR²³)(CH₂)_k- -(CH₂)_jC(H)(OR²³)(CH₂)_k- (in which latter three groups, the - (CH₂)_j- group is attached to the bispidine nitrogen atom, and which latter four groups are all optionally substituted by one or more OH groups);

B represents a single bond, C₁₋₄ alkylene, -(CH₂)_mN(R²⁴)-, (CH₂)_mS(O)_n-, -(CH₂)_mO- (in which three latter groups, the -(CH₂)_m- group is attached to the carbon atom bearing D and R⁴), -C(O)N(R²⁴)- (in which latter group, the -C(O)- group is attached to the carbon atom bearing D and R⁴), N(R²⁴)C(O)O(CH₂)_m- or -N(R²⁴)(CH₂)_m- (in which latter two groups, the N(R²⁴) group is attached to the carbon atom bearing D and R⁴);

j, k and m independently represent 0, 1, 2, 3 or 4;

n represents 0, 1 or 2;

R²³ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or C(O)R²⁵

R²⁴ represents H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

R²⁵ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Het³ or -(CH₂)_p-aryl (which latter two groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkyl and/or C₃₋₆ cycloalkoxy);

Het³ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

p represents 0, 1, 2, 3 or 4;


wherein alkyl groups that R¹, R⁴, R^{5a}, R^{5b}, R⁶, R⁷, R^{7a}, R^{7b}, R⁸, R¹⁰, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R^{18a}, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and D may represent, and with

which R^1 , R^7 , R^8 , R^{11} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} and R^{25} may be substituted; and alkoxy groups and that R^6 may represent, and with which R^1 , R^7 , R^8 , R^{11} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} and R^{25} may be substituted, may be linear or, when there is a sufficient number (i.e. three) of carbon atoms, be branched and/or ~~cyclic~~ cycloalkyl or cycloalkoxy with carbon ranges as defined above, and wherein, when there is a sufficient number (i.e. four) of carbon atoms, such alkyl and alkoxy groups may also be part ~~cyclic/acyclic~~ cycloalkyl/acyclic or cycloalkoxy/acyclic with carbon ranges as defined above, and wherein such alkyl and alkoxy groups may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen and/or substituted by one or more fluoro groups; and

wherein alkylene groups that A and B may represent, and $-(CH_2)-$ containing groups that R^1 , R^7 , R^8 , R^{10} , R^{11} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{25} , A, B and D may include, may be linear or, when there is a sufficient number (i.e. two) of carbon atoms, be branched, and wherein such alkylene groups and $-(CH_2)-$ containing chains may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen,

provided that when R^{5a} and R^{5b} both represent H, then D does not represent H or OH.


25 (previously presented). A compound of formula XVII,



XVII

wherein R^4 , R^{5a} , R^{5b} , R^6 , A, B and D are as defined in Claim 1, provided that when R^{5a} and R^{5b} both represent H, then D does not represent H or OH.

26 (currently amended). A process for the preparation of a compound of formula VIII,



VIII

wherein R^1 represents C_{1-12} alkyl, C_{3-12} cycloalkyl, $-(CH_2)_a$ -aryl, or $(CH_2)_a$ Het¹ (all of which are optionally substituted by one or more substituents selected from -OH, halo, cyano, nitro, C_{1-4} alkyl, C_{3-4} cycloalkyl and/or C_{1-4} alkoxy or C_{3-4} cycloalkoxy);

a represents 0, 1, 2, 3, or 4;

Het¹ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

X represents O or S;

R^{5a} and R^{5b} independently represent H, C_{1-3} alkyl or C_3 cycloalkoxy;

R^4 represents H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

D represents H, -OH, or $-(CH_2)_cN(R^{10})(R^{11})$;

c represents 0, 1, 2, 3 or 4;

R^{10} represents H, C_{1-6} alkyl, C_{3-6} cycloalkyl, $-(CH_2)_d$ -aryl, $-C(NH)NH_2$, $-S(O)_2R^{13}$, $-[C(O)]_eN(R^{14})(R^{15})$, $-C(O)R^{16}$ or $-C(O)OR^{17}$;

e represents 1 or 2;

R^{11} represents H, C_{1-6} alkyl, $-C(O)R^{18}$ or $-(CH_2)_f$ -aryl (which latter group is optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} independently represent H, C_{1-6} alkyl, C_{3-6} cycloalkyl, Het² or $-(CH_2)_g$ -aryl (which latter three groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

R^{13} represents C_{1-6} alkyl, C_{3-6} cycloalkyl, aryl or $-(CH_2)_h$ -aryl (all of which are all optionally substituted by one or more substituents chosen from halo, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkyl and/or C_{3-6} cycloalkoxy);

d, f, g and h independently represent 0, 1, 2, 3 or 4;

Het² represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;

R^6 represents one or more optional substituents selected from -OH, cyano, halo, amino, nitro, C_{1-6} alkyl (optionally terminated by $-N(H)C(O)OR^{18a}$), C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{3-6} cycloalkoxy, $-C(O)N(H)R^{19}$, $-NHC(O)N(H)R^{20}$, $-N(H)S(O)_2R^{21}$ and/or $-OS(O)_2R^{22}$;

R^{19} and R^{20} independently represent H, C_{1-6} alkyl or C_{3-6} cycloalkyl;

R^{18a} , R^{21} and R^{22} independently represent C_{1-6} alkyl or C_{3-6} cycloalkyl;

A represents a single bond, C_{1-6} alkylene, $-N(R^{23})(CH_2)_j-$, $-O(CH_2)_j-$ or $-(CH_2)_jC(H)(OR^{23})(CH_2)_k-$ $-(CH_2)_jC(H)(OR^{23})(CH_2)_k-$ (in which latter three groups, the $-(CH_2)_j-$ group is attached to the bispidine nitrogen atom, and which latter four groups are all optionally substituted by one or more OH groups);

B represents a single bond, C_{1-4} alkylene, $-(CH_2)_mN(R^{24})-$, $(CH_2)_mS(O)_n-$, $-(CH_2)_mO-$ (in which three latter groups, the $-(CH_2)_m-$ group is attached to the carbon atom bearing D and R^4), $-C(O)N(R^{24})-$ (in which latter group, the $-C(O)-$ group is attached to the carbon atom bearing D and R^4), $N(R^{24})C(O)O(CH_2)_m-$ or $-N(R^{24})(CH_2)_m-$ (in which latter two groups, the $N(R^{24})$ group is attached to the carbon atom bearing D and R^4);

j, k and m independently represent 0, 1, 2, 3 or 4;

n represents 0, 1 or 2;

R²³ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or C(O)R²⁵

R²⁴ represents H, C₁₋₆ alkyl or C₃₋₆ cycloalkyl;

R²⁵ represents H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Het³ or -(CH₂)_p-aryl (which latter two groups are optionally substituted by one or more substituents selected from -OH, cyano, halo, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₃₋₆ cycloalkyl and/or C₃₋₆ cycloalkoxy);

Het³ represents a five to ten-membered heterocyclic ring containing one or more heteroatoms selected from oxygen, nitrogen and/or sulfur, and which also optionally includes one or more =O substituents;


p represents 0, 1, 2, 3 or 4;

wherein alkyl groups that R¹, R⁴, R^{5a}, R^{5b}, R⁶, R⁷, R^{7a}, R^{7b}, R⁸, R¹⁰, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R^{18a}, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and D may represent, and with which R¹, R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted; and alkoxy groups and that R⁶ may represent, and with which R¹, R⁷, R⁸, R¹¹, R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸ and R²⁵ may be substituted, may be linear or, when there is a sufficient number (i.e. three) of carbon atoms, be branched and/or ~~cyclic~~ cycloalkyl or cycloalkoxy with carbon ranges as defined above, and wherein, when there is a sufficient number (i.e. four) of carbon atoms, such alkyl and alkoxy groups may also be part ~~cyclic/acyclic~~ cycloalkylacyclic or cycloalkoxy/acyclic with carbon ranges as defined above, and wherein such alkyl and alkoxy groups may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen and/or substituted by one or more fluoro groups; and

wherein alkylene groups that A and B may represent, and $-(CH_2)-$ containing groups that R^1 , R^7 , R^8 , R^{10} , R^{11} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{25} , A, B and D may include, may be linear or, when there is a sufficient number (i.e. two) of carbon atoms, be branched, and wherein such alkylene groups and $-(CH_2)-$ containing chains may ~~also be saturated or~~, when there is a sufficient number (i.e. two) of carbon atoms, be unsaturated and/or interrupted by oxygen,


provided that when R^{5a} and R^{5b} both represent H, then D does not represent H or OH, or

a compound of formula XVII,




XVII

wherein R^4 , R^{5a} , R^{5b} , R^6 , A, B and D are as defined in Claim 1, provided that when R^{5a} and R^{5b} both represent H, then D does not represent H or OH, which comprises reaction of a compound of formula XXIX,



XXIX

wherein R^2 represents H or $-C(O)XR^1$ and R^1 , R^{5a} , R^{5b} and X are as defined in Claim 1 with a compound of formula XXX,



XXX

wherein R^4 , R^6 , A, B and D are as defined in Claim 1, in the presence of a formaldehyde.

27 (previously presented). A method as claimed in Claim 20, wherein the arrhythmia is an atrial or a ventricular arrhythmia.